

Type 1 Diabetes May Have an Impact on the Developing Brain in Children

Nemours Researchers part of the Diabetes Research in Children Network Present Pivotal Research at American Diabetes Association Scientific Sessions

SAN FRANCISCO (June 9, 2019) – A study co-led by Dr. Nelly Mauras at Nemours Children’s Health System in Jacksonville, Florida found that children with type 1 diabetes (T1D) have slower growth in brain areas associated with mild cognitive deficits compared to children without T1D. The study, presented today at the American Diabetes Association’s® (ADA’s) 79th Scientific Sessions®, found significant differences in total brain and regional gray and white matter growth based on a series of three structural magnetic resonance imaging (MRI) studies.

“Even with new insulin and technologies that can significantly improve care, children with Type 1 Diabetes are still exposed to significant swings in sugar control, creating potential risks to the developing brain,” said Nelly Mauras, MD.

“Even with new insulin and technologies that can significantly improve care, children with Type 1 Diabetes are still exposed to significant swings in sugar control, creating potential risks to the developing brain,” said Nelly Mauras, MD, co-principal investigator of the study, chief of the division of endocrinology, diabetes & metabolism at the Nemours Children’s Health System in Jacksonville, Florida, and professor of pediatrics at the Mayo College of Medicine. “Understanding the early effects of blood sugar control on brain development is a necessary step towards developing strategies for reducing these risks and the public health implications of diabetes-related cognitive dysfunction later on in life.”

As part of a multi-site study of the Diabetes Research in Children Network (DirecNet), researchers aimed to determine the extent to which glycemic exposure adversely impacts the developing brain in children with early-onset T1D. The study enrolled 138 children with T1D with a median age of seven years. The participants had a disease duration on average of 2.4 years at the beginning of the study. MRIs were performed at three time points (baseline visit, 18 months and approximately 2.9 years after the second visit) to measure gray and white matter volumes in key brain regions. Total cumulative hyperglycemic exposure was determined using lifetime blood sugar, using hemoglobin A1c (HbA1c) values from the time of diagnosis. Researchers compared the MRI results of T1D participants to those of a control group of 66 age-matched children who did not have diabetes.

Researchers found that the group with T1D had slower growth of total cortical and subcortical gray and white matter than the control group at all time points. In particular, a set of metabolically active brain regions associated with other brain disorders, known as the “default mode network,” showed less growth in the T1D group compared to the control group. These regions of slower growth were associated with higher lifetime blood sugar, as measured by HbA1c values.

“Ongoing research is investigating whether diligent maintenance of blood sugar levels in the normal range through advanced diabetes technologies can impact these findings to reduce the risk for cognitive dysfunction,” said Mauras.

The DirecNet includes Nemours Children’s Health System JAX, Stanford University, University of Iowa, Washington University in St Louis, and Yale University. The research is supported through funding from the National Institutes of Health’s Special Type 1 Diabetes Funds and the Eunice Kennedy Shriver National Institute of Child Health and Human Development. Dr. Mauras and her co-principal investigator, Allan Reiss, MD, Howard C. Robbins Professor of Psychiatry and Behavioral Sciences and Professor of Radiology at Stanford University School of Medicine, will present these findings in a press briefing on Sunday, June 9, 2019 from 12-1 p.m. PT at the American Diabetes Association’s 79th Scientific Sessions in San Francisco.

###

[Nemours](#) is an internationally recognized children's health system that owns and two free-standing children's hospitals: the Nemours/Alfred I. duPont Hospital for Children in Wilmington, Del., and Nemours Children's Hospital in Orlando, Fla., along with outpatient facilities in five states, delivering pediatric primary, specialty and urgent care. Nemours also powers the world's most-visited website for information on the health of children and teens, [KidsHealth.org](#), and offers on-demand, online video patient visits through Nemours [CareConnect](#). [Nemours ReadingBrightstart.org](#) is a program dedicated to preventing reading failure in young children, grounded in Nemours' understanding that child health and learning are inextricably linked, and that reading level is a strong predictor of adult health.

Established as [The Nemours Foundation](#) through the legacy and philanthropy of Alfred I. duPont, Nemours provides pediatric clinical care, research, education, advocacy and prevention programs to families in the communities it serves.

Additional assets available online: [Photos \(1\)](#)

<https://nemours.mediaroom.com/2019-June-09-Diabetes-Impact-Developing-Brain>